

A SIPHONOTID MILLIPEDE (*RHINOTUS*) AS THE SOURCE OF SPIROPYRROLIZIDINE OXIMES OF DENDROBATID FROGS

R. A. SAPORITO,¹ M. A. DONNELLY,¹ R. L. HOFFMAN,²
H. M. GARRAFFO,³ and J. W. DALY^{3*}

¹Department of Biological Sciences, Florida International University, Miami, FL, 33199

²Virginia Museum of Natural History, 1001 Douglas Avenue, Martinsville, VA, 24112

³Laboratory of Bioorganic Chemistry, National Institute of Diabetes and Digestive
and Kidney Diseases, NIH, DHHS, Bethesda, MD, 20892-0820

(Received October 13, 2003; accepted October 10, 2002)

Abstract—Poison frogs of the neotropical family Dendrobatidae contain a wide variety of lipophilic alkaloids, which are accumulated from alkaloid-containing arthropods. A small millipede, *Rhinotus purpureus* (Siphonotidae), occurs microsympatrically with the dendrobatid frog *Dendrobates pumilio* on Isla Bastimentos, Bocas del Toro Province, Panamá. Methanol extracts of this millipede contain the spiropyrrolizidine *O*-methyloxime **236**, an alkaloid previously known only from skin extracts of poison frogs, including populations of *D. pumilio*. Thus, *R. purpureus* represents a likely dietary source of such alkaloids in dendrobatid frogs.

Key Words—Alkaloids, arthropods, frogs, dendrobatids, millipedes, and pyrrolizidines.

INTRODUCTION

Millipedes produce a wide range of allomones, which presumably act to deter potential predators. Millipede allomones include benzoquinones, phenols, benzaldehyde/hydrogen cyanide, and alkaloids (Eisner et al., 1978 and references therein). Three groups of alkaloids are known from millipedes as follows: (1) the glomerins (Ia & b) from millipedes of the genera *Glomeris* and *Loboglomeris* (Order: Glomerida, Family: Glomeridae) (Schildknecht et al., 1967 and reference therein), (2) the terpenoid alkaloids polyzonimine (II) and nitropolyzonamine (III) from the

* To whom correspondence should be addressed. E-mail: jdaly@nih.gov




FIG. 1. Millipede alkaloids.

millipede *Polyzonium rosalbum* [now *Petaserpes crytocephalus*] (Order: Polyzoniida, Family: Polyzoniidae) (Meinwald et al., 1975 and reference therein), and (3) the terpenoid alkaloid buzonamine (IV) from a millipede of the genus *Buzonium* (Order: Polyzoniida, Family: Polyzoniidae) (Wood et al., 2000) (Figure 1). Certain poison frogs/toads of the families Dendrobatidae, Mantellidae, and Bufonidae contain spiropyrrolizidine alkaloids (Figure 1; Va & b and VI) (Daly et al., 1994 and references therein) with carbon skeletons identical to those of the millipede alkaloid nitropolyzonamine. These spiropyrrolizidine oximes have been assigned 'bold type' code names (**222**, **236**, and **252A**), which consist of a boldfaced number corresponding to the nominal mass and a boldfaced letter for identification of individual alkaloids. Such frogs appear to obtain their skin alkaloids from dietary sources (Daly et al., 1994, 2000), and, therefore, it seemed likely that a polyzoniid millipede would prove to be a dietary source of spiropyrrolizidine oximes.

Spiropyrrolizidine **236** and nitropolyzonamine have been detected in extracts of leaf-litter arthropods collected with Berlese Funnels from Cerro Ancón, Panamá (Daly et al., 1994) and in mixed leaf-litter samples from Isla Bastimentos, Panamá (Daly et al., 2002), however, the specific arthropod source was not determined. In an effort to taxonomically identify arthropod sources for the frog skin alkaloids, a series of leaf-litter arthropod samples, separated by taxa (including millipedes), were again collected and analyzed from Isla Bastimentos, Panamá. From these collections, the spiropyrrolizidine oxime **236** was detected in samples containing only millipedes. We now report on the identification of a small millipede, *Rhinotus purpureus* (Pocock, 1894) (Order: Polyzoniida, Family: Siphonotidae), that

contains the spiropyrrolizidine oxime **236** and, thus, represents a dietary source for the same alkaloid found in the skin extracts of the dendrobatid frogs (*Dendrobates pumilio*) from the same location on Isla Bastimentos. In addition, trace amounts of nitropolyzonamine were detected in the millipede extract. Based on stomach content analysis, the diet of *D. pumilio* consists of a variety of small leaf-litter arthropods, including small millipedes (Donnelly unpublished data; Saporito et al., unpublished data).

METHODS AND MATERIALS

A variety of arthropods were collected from the leaf-litter and separated by taxa at several different sites on Isla Bastimentos, Bocas del Toro Province, Panamá. Arthropod collections were made from February 2–8, 2003, from the following 8 locations: Site 1: N 9°21.618'; W 82°12.074'. Site 2: N 9°21.250'; W 82°12.519'. Site 3: N 9°21.169'; W 82°12.627'. Site 4: N 9°21.123'; W 82°12.620'. Site 5: N 9°20.996'; W 82°12.726'. Site 6: N 9°20.364'; W 82°10.807'. Site 7: N 9°21.021'; W 82°12.704'. Site 8: N 9°20.490'; W 82°10.486'. Several of these sites were at the same location where mixed arthropod collections were previously collected (Daly et al., 2002). All arthropod specimens were collected with forceps, from leaf-litter placed on white cloth, and placed in plastic 150 ml vials according to individual taxa. Only small arthropods (less than 10 mm), suitable as prey for the dendrobatid frog *D. pumilio* (average snout-vent length 20 mm), were collected from these sites.

Gas chromatographic mass spectral analyses (GC-MS) were conducted with a Finnigan GCQ instrument with a 25 m × 0.25 mm i.d. Rtx-5 Amine fused silica column, using a temperature program from 100 to 280°C at a rate of 10°C/min. Arthropod extracts were screened for the presence of alkaloids by injecting a 4 μ l mixture, which consisted of 1 μ l portions from each of 4 different vials (arthropod samples). All mixtures were based on specific taxonomic groups (i.e., millipede samples were only combined with other millipede samples, etc.). If an alkaloid was detected in a mixture, then each of the four vials was analyzed separately to determine the source of the alkaloid. Identification of alkaloids was based on gas chromatographic mass spectra and comparison to authentic spiropyrrolizidine **236** or nitropolyzonamine.

RESULTS AND DISCUSSION

The spiropyrrolizidine alkaloid **236** was detected in two separate millipede samples, both of which were from site 7 located on Isla Bastimentos. Both samples contained one individual of *Rhinotus purpureus* (Pocock, 1894). In addition, trace amounts of nitropolyzonamine were also found in these samples.

Alkaloids were not detected in extracts from the following other millipede samples: (1) Order: Polyxenida; 1 individual. (2) Order: Stemmiulidae; Family: Stemmiulidae; Species: *Prostemmiulus* cf. *cinnatus* (Loomis); 3 individuals. (3) Order: Spirobolida; Family: Spirobolellidae; Species: *Spirobolellus trifasciatus* (Loomis); 3 individuals. (4) Order: Polydesmida; Family: Aphelidesmidae; Genus: *Aphelidesmus*; 1 adult and 1 immature. (5) Order: Polydesmida; Family: Chelodesmidae; Genus: *Trichomorpha*; 1 individual. (6) Order: Polydesmida; Family: Cyrtodesmidae; Genus: *Cyrtodesmus*; 1 individual. (7) Order: Polydesmida; Family: Fuhrmannodesmidae; 2 individuals. (8) Order: Polydesmida; 3 immature individuals.

Spiropyrrrolizidine **236** was not detected in any of the other arthropod extracts examined in this study, however, a decahydroquinoline and a 5,6,8-trisubstituted indolizidine were detected in certain ant extracts. These results will be reported separately. Decahydroquinolines have been reported from myrmicine ants (Daly et al., 2000, 2002 and references therein). However, until now, there have been no known reports of branched chain izidine alkaloids from ant extracts, although a branched chain 5,8-disubstituted indolizidine was detected in extracts from mixed arthropod samples from Isla Bastimentos (Daly et al., 2002). In the present study, the frogs from the site at which the millipedes had the spiropyrrrolizidine **236** were also shown to have that alkaloid in skin extracts (Daly et al., 2002 and unpublished data). Nitropolyzonamine was not detected in the skin extracts from *D. pumilio* on Isla Bastimentos. However, nitropolyzonamine, as well as the spiropyrrrolizidine oximes **236** and **252B** (not shown, see below) have been detected as trace amounts in *Dendrobates auratus* from Cerro Ancon, Panamá (Daly et al., 1994).

The spiropyrrrolizidine oxime **236** has been detected in frog skin extracts from several species of dendrobatid frogs from a variety of locations throughout Central and South America (Table 1). From these locations, it is interesting to note the absence of **236** from some of the frog populations in Panamá during initial visits (Table 1). The appearance of **236** in later visits to the same Panamanian sites may suggest an introduction and, therefore, availability of *R. purpureus* to frogs within these populations.

The millipede *R. purpureus* has a relatively wide distribution throughout the tropics; therefore, it seems probable that other *Rhinotus* species may serve as a source for spiropyrrrolizidine oximes found in dendrobatid frogs of South America. Spiropyrrrolizidine oxime **236** has been detected from an Argentinean bufonid toad (Garraffo et al., 1993) and from Madagascan mantellid frogs (Daly et al., 1996). In addition, one extract from an Australian myobatrachid frog had spiropyrrrolizidine oxime **252B**, an isomer of **252A** in which the hydroxyl group is in a different position (Daly et al., 1990). It is likely that such spiropyrrrolizidine oximes will be found to have a deterrent effect on potential millipede predators,

TABLE 1. OCCURRENCE OF SPIROPYRROLIZIDINE **236** IN FROG/TOAD SKIN EXTRACTS: AN INDICATION OF AVAILABILITY OF *Rhinotus* MILLIPEDES?

| Family: <i>Species</i> | Location | Year | 236 |
|-----------------------------------|---|-------------------------------------|--------------------|
| Dendrobatidae: | | | |
| <i>Dendrobates pumilio</i> | Rumpala, Bocas Province, Panamá | 1981 | Present |
| | Isla Bastimentos, Bocas Province, Panamá | 1972 | Absent† |
| | | 1980, 1981, 1987, 1993, 2002* | Present |
| | La Gruta, Isla Colon, Bocas Province, Panamá | 1983, 1986 1992 | Absent† Present |
| | Río Sand Box, Limón Province, Costa Rica | 1989 | Present |
| | Río Sarapiquí, Heredia Province, Costa Rica | 1989 | Present |
| | 28 Millas, Limón Province, Costa Rica | 1990, 1995 | Present |
| | Siquirres, Limón Province, Costa Rica | 1995 | Present |
| | Pocora, Limón Province, Costa Rica | 1995 | Present |
| | <i>Dendrobates auratus</i> | Isla Tobago, Panamá | 1974 |
| | | 1993 | Present |
| | Cerro Ancón, Panamá | 1993 | Present |
| <i>Dendrobates tinctorius</i> | Pet Trade | 1985 | Present |
| <i>Epipedobates macero</i> | Parque Nacional Manu, Perú | 1989 | Present |
| <i>Epipedobates cf. pictus</i> | Parque Nacional Manu, Perú | 1989 | Present |
| <i>Epipedobates pulchripectus</i> | Serra de Navio, Amapa, Brazil | 1992 | Present |
| <i>Epipedobates tricolor</i> | Pasaje, El Oro, Ecuador | 1987 | Present |
| <i>Epipedobates trivittatus</i> | Pet Trade | 1985 | Present |
| Mantellidae: | | | |
| <i>Mantella baroni</i> | Sahavondrana, Madagascar | 1993 | Present |
| <i>Mantella betsileo</i> | Mahambo, Madagascar | 1998 | Present |
| Bufonidae: | | | |
| <i>Melanophryniscus stelzneri</i> | Tanti, Cordoba, Argentina | 1989 | Present |

*see Daly et al. 2002.

†not detected.

such as ants, however, such studies have yet to be conducted. The oxime **236** does exert potent blocking action at mammalian nicotinic receptors, particularly those of the ganglionic subtype (Badio et al., 1996).

Acknowledgments—We would like to thank the Autoridad Nacional del Ambiente and the República de Panamá for permission to collect and export the arthropod specimens used in this study (Permit: SE/A-45-03). We thank the Smithsonian Tropical Research Institute (STRI), especially Orelis Arosemena and Maria Leone, for assistance in obtaining these permits. This work is based on research supported by the Environmental Protection Agency, the Explorers Club, and a Courtesy Associate appointment given by the National Institute of Diabetes and Digestive and Kidney Diseases. This paper is contribution number 67 to the program in Tropical Biology at Florida International University.

REFERENCES

- BADIO, B., SHI, D., SHIN, Y., HUTCHINSON, K. D., PADGETT, W. L., AND DALY, J. W. 1996. Spiropyrrolizidines: A new class of blockers of nicotinic receptors. *Biochem. Pharmacol.* 52:933–939.
- DALY, J. W., ANDRIAMAHARAVO, N. R., ANDRIANTSIFERANA, M., AND MYERS, C. W. 1996. Madagascar poison frogs (*Mantella*) and their skin alkaloids. *Am. Mus. Novitates* 3177:1–34.
- DALY, J. W., GARRAFFO, H. M., JAIN, P., SPANDE, T. F., SNELLING, R. R., JARAMILLO, C., AND RAND, A. S. 2000. Arthropod-frog connection: Decahydroquinoline and pyrrolizidine alkaloids common to microsymbiotic myrmicine ants and dendrobatid frogs. *J. Chem. Ecol.* 26:73–85.
- DALY, J. W., GARRAFFO, H. M., PANNELL, L. K., AND SPANDE, T. F. 1990. Alkaloids from Australian frogs (Myobatrachidae): Pseudophrynamines and pumiliotoxins. *J. Nat. Prod.* 53:407–421.
- DALY, J. W., GARRAFFO, M. H., SPANDE, T. F., JARAMILLO, C., AND RAND, S. A. 1994. Dietary source for skin alkaloids of poison frogs (Dendrobatidae)? *J. Chem. Ecol.* 20:943–955.
- DALY, J. W., KANEKO, T., WILHAM, J., GARRAFFO, M. H., SPANDE, T. F., ESPINOSA, A., AND DONNELLY, M. A. 2002. Bioactive alkaloids of frog skins: Combinatorial bioprospecting reveals that pumiliotoxins have an arthropod source. *Proc. Natl. Acad. Sci. U.S.A.* 99:13996–14001.
- EISNER, T., ALSOP, D., HICKS, K., AND MEINWALD, J. 1978. Defensive secretions of millipedes, pp. 41–72. *in*: Handbook of Experimental Pharmacology. Vol. 4. S. Bertini (ed.). Springer Verlag, Berlin, Germany.
- GARRAFFO, H. M., SPANDE, T. F., DALY, J. W., BALDESSARI, A., AND GROS, E. G. 1993. Alkaloids from bufonid toads (*Melanophryniscus*): Decahydroquinolines, pumiliotoxins and homopumiliotoxins, indolizidines, pyrrolizidines, and quinolizidines. *J. Nat. Prod.* 56:357–373.
- MEINWALD, J., SMOLANOFF, J., MCPHAIL, A. T., MILLER, R. W., EISNER, T., AND HICKS, K. 1975. Nitropolyzonamine: A spirocyclic nitro compound from the defensive glands of a millipede (*Polyzonium rosalbum*). *Tetrahedron Lett.* 28:2367–2370.
- SCHILDKNECHT, H., MASCHWITZ, U., AND WENNEIS, W. F. 1967. Neue Stoffe aus dem Wehrsekret der Diplopodengattung *Glomeris*. *Naturwissenschaften* 16:196–197.
- WOOD, W. F., HANKE, F. J., KUBO, I., CARROLL, J. A., AND CREWS, P. 2000. Buzonamine, a new alkaloid from the defensive secretions of the millipede, *Buzonium crassipes*. *Biochemi. Syst. Ecol.* 28:305–312.